

United Nations Educational Scientific and Cultural Organization  
and  
International Atomic Energy Agency  
INTERNATIONAL CENTRE FOR THEORETICAL PHYSICS

**MAXIMAL ELEMENTS OF SUPPORT AND COSUPPORT**

Siamak Yassemi<sup>1</sup>  
International Centre for Theoretical Physics, Trieste, Italy.

ABSTRACT

We introduce a set that is tightly closed to the set of the Jacobson radical of module (the intersection of all maximal elements in support). In the last section, it is proved that the set of zero divisors of a module is equal to the union of the maximal elements of the support of module if the module is finitely generated and injective.

MIRAMARE – TRIESTE

May 1997

---

<sup>1</sup>Permanent address: Department of Mathematics, University of Tehran, P.O. Box 13145-448, Tehran, Iran. E-mail address: [yassemi@rose.ipm.ac.ir](mailto:yassemi@rose.ipm.ac.ir)

## 0. Introduction

Throughout this note the ring  $R$  is commutative (not necessarily Noetherian) with non-zero identity. The notion of prime ideals is central to the commutative ring theory. The set  $\text{Spec}(R)$  of prime ideals of a ring  $R$  is a topological space, and the localization of rings with respect to this topology is an important technique for studying them. In addition, the maximal element of this set is very useful. There is a similar notion for modules that is the support of modules. The set of prime ideals  $\mathfrak{p}$  such that there exists a cyclic submodule  $N$  of  $M$  with  $\mathfrak{p} \supseteq \text{Ann}(N)$  is well-known to be the support of  $M$ , and is written  $\text{Supp}(M)$ . In [Y1] we have introduced the notion of cocyclic modules (that is, a submodule of  $E(R/\mathfrak{m})$ , the injective envelope of  $R/\mathfrak{m}$ , for some  $\mathfrak{m} \in \text{Max}R$ ) and it is used to define the notion of cosupport (the dual notion of support) over Noetherian rings. In this paper we define the notion of cosupport of modules over (not necessarily Noetherian) rings. We study the maximal element of support and cosupport of modules. Let  $J_R(M)$  be the Jacobson radical of the  $R$ -module  $M$  (intersection of all maximal elements of the support of  $M$ ). Let  $N_R(M)$  be the union of all maximal elements of the support of  $M$ . Then it is easy to see that  $J_R(M) \subseteq N_R(M)$  and there is equality if and only if the support of  $M$  has only one element. The set  $N'_R(M)$  is defined by

$$N'_R(M) = \{x \in M \mid x + N_R(M) \subseteq N_R(M)\}.$$

We show that  $J_R(M) \subseteq N'_R(M)$  and there is equality if the support of  $M$  has only finite elements. By an example we show that the inequality may be strict. In the last section we prove that for any finitely generated and injective  $R$ -module  $M$ , the set of zero divisors of  $M$  is equal to the set  $N_R(M)$ . As a corollary of this result we have that “if  $R$  is a self-injective ring then each non-unit element in  $R$  is a zero divisor in  $R$ ”.

## 1. Support of Modules

In this section we study the intersection and union of the maximal elements of the support of a module.

**Definition 1.1.** Let  $M$  be an  $R$ -module. The *support* of  $M$  is denoted by  $\text{Supp}(M)$  and

it is defined by

$$\text{Supp}(M) = \{\mathfrak{p} \in \text{Spec}(R) \mid \mathfrak{p} \supseteq \text{Ann}(N) \text{ for some cyclic submodule } N \text{ of } M\}.$$

The next lemma is a well-known result, cf. [M].

**Lemma 1.2.** *The following hold:*

- a)  $M \neq 0$  if and only if  $\text{Supp}(M) \neq \emptyset$ .
- b)  $\text{Supp}(M) \subseteq \text{Spec}(R/\text{Ann}(M))$ .
- c) If  $M$  is finite then we have equality in (b).

**Remark 1.3.** The inequality of (1.2b) may be strict, for example, if  $(R, \mathfrak{m})$  is a local ring and  $M = E(R/\mathfrak{m})$ , injective envelope of the field  $R/\mathfrak{m}$ , then  $\text{Ann}(M) = 0$  and so  $\text{Spec}(R/\text{Ann}(M)) = \text{Spec}(R)$ . On the other hand  $\text{Supp}(M) = \{\mathfrak{m}\}$ .

**Theorem 1.4.** *Let  $M$  be an  $R$ -module. Then*

$$\text{MaxSupp}(M) \subseteq \text{MaxSpec}(R/\text{Ann}(M))$$

where  $\text{MaxSupp}(M)$  is the set of all maximal elements in  $\text{Supp}(M)$ .

**Proof.** Set  $\mathfrak{p} \in \text{MaxSupp}(M)$ . Then by (1.2) we have that  $\mathfrak{p} \in \text{Spec}(R/\text{Ann}(M))$ . Choose  $\mathfrak{m} \in \text{MaxSpec}(R/\text{Ann}(M))$  such that  $\mathfrak{p} \subseteq \mathfrak{m}$ . By (1.1) it is easy to see that  $\mathfrak{m} \in \text{Supp}(M)$ . Since  $\mathfrak{p} \in \text{MaxSupp}(M)$  we have  $\mathfrak{p} = \mathfrak{m}$  and hence the assertion holds.  $\square$

**Remark 1.5.** The inequality in (1.4) may be strict. For example let  $R$  be an integral domain and  $\{\mathfrak{m}, \mathfrak{n}\} \subseteq \text{MaxSpec}(R)$ . Let  $M = E(R/\mathfrak{m})$ . Then  $\text{MaxSupp}(M) = \{\mathfrak{m}\}$  but  $\mathfrak{n} \in \text{MaxSpec}(R/\text{Ann}(M)) = \text{MaxSpec}(R)$ .

**Definition 1.6.** Let  $M$  be an  $R$ -module. The *Jacobson radical* of  $M$  is denoted by  $J_R(M)$  and it is the intersection of all elements in  $\text{MaxSupp}(M)$ . Also the union of all elements in  $\text{MaxSupp}(M)$  is denoted by  $N_R(M)$ .

**Lemma 1.7.** *Let  $M$  be an  $R$ -module. Then  $r \in J_R(M)$  if and only if  $1 + tr \notin N_R(M)$  for any  $t \in R$ .*

**Proof.** “if” Let  $\mathfrak{m} \in \text{MaxSupp}(M)$  such that  $r \notin \mathfrak{m}$ . Then  $\mathfrak{m} \in \text{Max}(R)$  and hence  $\mathfrak{m} + rR = R$ . Therefore, there exist  $x \in \mathfrak{m}$  and  $t \in R$  such that  $x + tr = 1$  and hence  $1 - rt \in N_R(M)$ , which is a contradiction.

“only if” Let  $t \in R$  such that  $1 + tr \in N_R(M)$ . Then there exists a maximal ideal  $\mathfrak{m} \in \text{MaxSupp}(M)$  such that  $1 + tr \in \mathfrak{m}$ . On the other hand  $tr \in \mathfrak{m}$ . Therefore  $1 \in \mathfrak{m}$ , which is a contradiction.  $\square$

**Definition 1.8.** The  $R$ -module  $M$  is said to be *local module* if  $|\text{MaxSupp}(M)| = 1$ . Also the  $R$ -module  $M$  is said to be *semi-local module* if  $|\text{MaxSupp}(M)| < \infty$ . Clearly, all non-zero modules over a semi-local (resp. local) ring is a semi-local (resp. local) module.

**Theorem 1.9.** *The following are equivalent:*

- i)  $M$  is a local module.
- ii)  $J_R(M) = N_R(M)$ .
- iii)  $N_R(M)$  is an ideal of  $R$ .

**Proof.** “(i $\Rightarrow$  ii)” and “(ii $\Rightarrow$  iii)” are obvious.

(iii $\Rightarrow$  i)” Since  $1 \notin N_R(M)$  we have  $N_R(M) \neq R$  and hence there exists  $\mathfrak{m} \in \text{MaxSpec}R$  such that  $N_R(M) \subseteq \mathfrak{m}$ . On the other hand  $\mathfrak{m} \subseteq N_R(M)$ . Therefore  $N_R(M) = \mathfrak{m}$  and hence  $\text{MaxSupp}(M) = \{\mathfrak{m}\}$ .  $\square$

**Definition 1.10.** Let  $M$  be an  $R$ -module. We define  $N'_R(M)$  by

$$N'_R(M) = \{x \in N_R(M) \mid x + N_R(M) \subseteq N_R(M)\}.$$

**Theorem 1.11.** *Let  $M$  be an  $R$ -module. Then the following hold:*

- a)  $J_R(M) \subseteq N'_R(M) \subseteq N_R(M)$
- b)  $J_R(M) = N'_R(M)$  if and only if  $N'_R(M)$  is an ideal of  $R$ .
- c) If  $M$  is a semi-local then  $J_R(M) = N'_R(M)$ .

**Proof.** “(a)” Set  $x \in J_R(M)$  and  $t \in N_R(M)$ . Then there exists  $\mathfrak{m} \in \text{MaxSupp}(M)$  such that  $t \in \mathfrak{m}$ . Since  $x \in \mathfrak{m}$  we have  $x + t \in \mathfrak{m}$  and hence  $x + t \in N_R(M)$ . Thus  $J_R(M) \subseteq N'_R(M)$ .

“(b)” The ‘Only if’ part is obvious. For the ‘If’ part, set  $x \in N'_R(M)$  and  $t \in R$ . Since  $N'_R(M)$  is an ideal of  $R$  we have  $tx \in N'_R(M)$ . We claim that  $1 + tx \notin N_R(M)$ . In the other case if  $1 + tx \in N_R(M)$  then  $1 \in N_R(M)$ , which is a contradiction. Therefore  $x \in J_R(M)$ .

“(c)” Let  $\text{MaxSupp}(M) = \{\mathfrak{m}_1, \mathfrak{m}_2, \dots, \mathfrak{m}_t\}$ . Suppose  $x \in N'_R(M)$ . Then there exists  $1 \leq r \leq t$  such that  $x \in \bigcap_{i=1}^r \mathfrak{m}_i$  and  $x \notin \bigcup_{i=r+1}^t \mathfrak{m}_i$ . We claim that  $r = t$ . In the other case by the prime avoidance theorem we have  $\bigcap_{i=r+1}^t \mathfrak{m}_i \not\subseteq \bigcup_{i=1}^r \mathfrak{m}_i$  and hence there exists  $y \in \bigcap_{i=r+1}^t \mathfrak{m}_i \setminus \bigcup_{i=1}^r \mathfrak{m}_i$ . Since  $y \in N_R(M)$  we have  $x + y \in N_R(M)$ . On the other hand  $x + y \notin \mathfrak{m}_i$  for each  $1 \leq i \leq t$ , which is a contradiction.  $\square$

**Remark 1.12.** The inequalities in 1.11 (a) may be strict. For the inequality in the right-hand side let  $M$  be a semi-local module but not local then  $J_R(M) = N'_R(M) \not\subseteq N_R(M)$ . For the inequality in the left-hand side, let  $(D, \mathfrak{m})$  be a local regular ring that is not a field. Then  $J_D(D) = \mathfrak{m} \neq 0$ . It is easy to see that  $J_{D[x]}(D[x]) = 0$  and

$$N_{D[x]}(D[x]) = N_D(D) \cup \{g \in D[x] \mid \deg(g) \geq 1\}.$$

Now we show that  $N'_D(D) \subseteq N'_{D[x]}(D[x])$ . Assume that  $a \in N'_D(D)$  then  $a \in N_D(D)$  and hence  $a \in N_{D[x]}(D[x])$ . Let  $f \in N_{D[x]}(D[x])$ . Then we have two cases:

- (i) “ $f \in N_D(D)$ ” In this case we have  $a + f \in N_D(D)$  and hence  $a + f \in N_{D[x]}(D[x])$ .
- (ii) “ $f \in D[x]$  with  $\deg f \geq 1$ ” Let  $f = \sum_{i=0}^n a_i x^i$  and let  $a_n \neq 0$ . Then  $\deg a + f \geq 1$  and hence  $a + f \in N_{D[x]}(D[x])$ . Therefore  $a \in N'_{D[x]}(D[x])$  and so  $N'_D(D) \subseteq N'_{D[x]}(D[x])$ .

Since  $0 \neq J_D(D) \subseteq N'_D(D) \subseteq N'_{D[x]}(D[x])$  we have that  $N'_{D[x]}(D[x]) \neq 0$ . On the other hand  $J_{D[x]}(D[x]) = 0$ .

By using the next lemma we can put  $N'_R(M)$  instead of  $J(M)$  in the Nakayama lemma and in the Krull’s intersection theorem.

**Lemma 1.13.** *Let  $M$  be an  $R$ -module and let  $\mathfrak{a}$  be an ideal of  $R$ . Then  $\mathfrak{a} \subseteq N'_R(M)$  if and only if  $\mathfrak{a} \subseteq J_R(M)$ .*

**Proof.** Let  $x \in \mathfrak{a}$  and  $r \in R$ . Then  $rx \in \mathfrak{a}$  and hence  $1 + rx \notin N'_R(M)$ . Therefore  $x \in J_R(M)$  by (1.7).  $\square$

## 2. Cosupport of modules

In this section we study the intersection and union of the maximal elements of the cosupport of a module.

**Definition 2.1** (see[Y1]). An  $R$ -module  $L$  is said to be cocyclic, if  $L$  is a submodule of  $E(R/\mathfrak{m})$ , the injective envelope of  $R/\mathfrak{m}$ , for some  $\mathfrak{m} \in \text{Max}R$ .

**Definition 2.2.** Let  $M$  be an  $R$ -module. The *cosupport* of  $M$  is denoted by  $\text{Cosupp}(M)$  and it is defined by

$$\text{Cosupp}(M) = \{\mathfrak{p} \in \text{Spec}(R) \mid \mathfrak{p} \supseteq \text{Ann}(L) \text{ for some cocyclic homomorphic image } L \text{ of } M\}.$$

**Lemma 2.3.** *The following hold:*

- a)  $M \neq 0$  if and only if  $\text{Cosupp}(M) \neq \emptyset$ .
- b)  $\text{Cosupp}(M) \subseteq \text{Spec}(R/\text{Ann}(M))$ .
- c) If  $M$  is finitely cogenerated then we have equality in (b).

**Proof.** “(a)” Use [Y2; 2.10 and 3.7].

“(b)” Let  $\mathfrak{p} \in \text{Cosupp}(M)$ . Then there exists a cocyclic homomorphic image  $L$  of  $M$  such that  $\mathfrak{p} \supseteq \text{Ann}(L)$ . Therefore  $\mathfrak{p} \supseteq \text{Ann}(M)$ .

“(c)” Since  $M$  is a finitely cogenerated  $R$ -module, there exists simple modules  $S_1, S_2, \dots, S_n$  such that  $E(M) = E(S_1) \oplus E(S_2) \oplus \dots \oplus E(S_n)$ . If  $\mathfrak{p} \in \text{Spec}(R/\text{Ann}(M))$ , then  $\mathfrak{p} \supseteq \bigcap_{i=1}^n \text{Ann}(E(S_i))$ . It follows that  $\mathfrak{p} \supseteq \text{Ann}(E(S_i))$  for some  $1 \leq i \leq n$ , thus  $\mathfrak{p} \in \text{Cosupp}(M)$ . Now the assertion follows from (b).  $\square$

**Theorem 2.4.** *Let  $M$  be an  $R$ -module. Then*

$$\text{MaxCosupp}(M) \subseteq \text{MaxSpec}(R/\text{Ann}(M))$$

where  $\text{MaxCosupp}(M)$  is the set of maximal element in  $\text{Cosupp}(M)$ .

**Proof.** Set  $\mathfrak{p} \in \text{MaxCosupp}(M)$ . Then by (2.3) we have that  $\mathfrak{p} \in \text{Spec}(R/\text{Ann}(M))$ . Choose  $\mathfrak{m} \in \text{MaxSpec}(R/\text{Ann}(M))$  such that  $\mathfrak{p} \subseteq \mathfrak{m}$ . It is easy to see that  $\mathfrak{m} \in \text{Cosupp}(M)$ . Since  $\mathfrak{p} \in \text{MaxCosupp}(M)$  we have that  $\mathfrak{p} = \mathfrak{m}$  and hence the assertion holds.  $\square$

**Theorem 2.5.** *Let  $M$  be an  $R$ -module. Then the following hold:*

- a)  $\text{MaxSupp}(M) \subseteq \text{MaxCosupp}(M)$
- b) *We have equality in (a) if  $M$  is finitely generated.*

Note: The inequality in (a) may be strict.

**Proof.** Use [Y1; 2.11].  $\square$

**Definition 2.6.** Let  $M$  be an  $R$ -module. The *Jacobson coradical* of  $M$  is denoted by  $C_R(M)$  and it is the intersection of all elements in  $\text{MaxCosupp}(M)$ . Also, the union of all elements in  $\text{MaxCosupp}(M)$  is denoted by  $U_R(M)$ .

**Lemma 2.7.** *Let  $M$  be an  $R$ -module. Then  $r \in C_R(M)$  if and only if  $1 + tr \notin U_R(M)$  for any  $t \in R$ .*

**Proof.** “if” Let  $\mathfrak{m} \in \text{MaxCosupp}(M)$  such that  $r \notin \mathfrak{m}$ . Then  $\mathfrak{m} \in \text{Max}(R)$  and hence  $\mathfrak{m} + rR = R$ . Therefore there exist  $x \in \mathfrak{m}$  and  $t \in R$  such that  $x + tr = 1$  and hence  $1 - rt \in U_R(M)$ , which is a contradiction.

“only if” Let  $t \in R$  such that  $1 + tr \in U_R(M)$ . Then there exists a maximal ideal  $\mathfrak{m} \in \text{MaxCosupp}(M)$  such that  $1 + tr \in \mathfrak{m}$ . On the other hand  $tr \in \mathfrak{m}$ . Therefore  $1 \in \mathfrak{m}$ , which is a contradiction.  $\square$

**Definition 2.8.** The  $R$ -module  $M$  is said to be *colocal module* if  $|\text{MaxCosupp}(M)| = 1$ . Also the  $R$ -module  $M$  is said to be *semi-colocal module* if  $|\text{MaxCosupp}(M)| < \infty$ . Clearly, all non-zero modules over a semi-local (resp. local) ring is a semi-local (resp. local) module. Also if  $M$  is a colocal (resp. semi-colocal) module then  $M$  is a local (resp. semi-local) module.

**Theorem 2.9.** *The following are equivalent:*

- i)  *$M$  is a colocal module.*

ii)  $C_R(M) = U_R(M)$ .

iii)  $U_R(M)$  is an ideal of  $R$ .

**Proof.** “(i $\Rightarrow$  ii)” and “(ii $\Rightarrow$  iii)” are obvious.

“(iii $\Rightarrow$  i)” Since  $1 \notin U_R(M)$  we have  $U_R(M) \neq R$  and hence there exists  $\mathfrak{m} \in \text{MaxSpec}R$  such that  $U_R(M) \subseteq \mathfrak{m}$ . On the other hand  $\mathfrak{m} \subseteq U_R(M)$ . Therefore  $U_R(M) = \mathfrak{m}$  and hence  $\text{MaxCosupp}(M) = \{\mathfrak{m}\}$ .  $\square$

**Definition 2.10.** Let  $M$  be an  $R$ -module. We define  $U'_R(M)$  by

$$U'_R(M) = \{x \in U_R(M) \mid x + U_R(M) \subseteq U_R(M)\}.$$

**Theorem 2.11.** Let  $M$  be an  $R$ -module. Then the following hold:

a)  $C_R(M) \subseteq U'_R(M) \subseteq U_R(M)$

b)  $C_R(M) = U'_R(M)$  if and only if  $U'_R(M)$  is an ideal of  $R$ .

c) If  $M$  is a semi-colocal then  $C_R(M) = U'_R(M)$ .

**Proof.** “(a)” Set  $x \in C_R(M)$  and  $t \in U_R(M)$ . Then there exists  $\mathfrak{m} \in \text{MaxCosupp}(M)$  such that  $t \in \mathfrak{m}$ . Since  $x \in \mathfrak{m}$  we have  $x + t \in \mathfrak{m}$  and hence  $x + t \in U_R(M)$ . Thus  $C_R(M) \subseteq U'_R(M)$ .

“(b)” The ‘Only if’ part is obvious. For the ‘If’ part, set  $x \in U'_R(M)$  and  $t \in R$ . Since  $U'_R(M)$  is an ideal of  $R$  we have  $tx \in U'_R(M)$ . We claim that  $1 + tx \notin U_R(M)$ . In the other case if  $1 + tx \in U_R(M)$  then  $1 \in U_R(M)$ , which is a contradiction. Therefore  $x \in C_R(M)$ .

“(c)” Let  $\text{MaxCosupp}(M) = \{\mathfrak{m}_1, \mathfrak{m}_2, \dots, \mathfrak{m}_t\}$ . Suppose  $x \in U'_R(M)$ . Then there exists  $1 \leq r \leq t$  such that  $x \in \bigcap_{i=1}^r \mathfrak{m}_i$  and  $x \notin \bigcup_{i=r+1}^t \mathfrak{m}_i$ . We claim that  $r = t$ . In the other case by the prime avoidance theorem we have  $\bigcap_{i=r+1}^t \mathfrak{m}_i \not\subseteq \bigcup_{i=1}^r \mathfrak{m}_i$  and hence there exists  $y \in \bigcap_{i=r+1}^t \mathfrak{m}_i \setminus \bigcup_{i=1}^r \mathfrak{m}_i$ . Since  $y \in U_R(M)$  we have  $x + y \in U_R(M)$ . On the other hand  $x + y \notin \mathfrak{m}_i$  for each  $1 \leq i \leq t$ , which is a contradiction.  $\square$

### 3. Injective and flat modules

Recall that the set of zero divisors of  $M$ ,  $Z_R(M)$ , is defined by

$$Z_R(M) = \{a \in R \mid M \xrightarrow{a} M \text{ is not injective}\}$$

Now we bring the dual notion of  $Z_R(M)$ .

**Definition 3.1**(see [Y1]). For the  $R$ -module  $M$  the subset  $W_R(M)$  of  $R$  is defined by

$$W_R(M) = \{a \in R \mid M \xrightarrow{a} M \text{ is not surjective}\}.$$

**Lemma 3.2.** *Let  $M$  be an  $R$ -module. Then the following hold;*

- a)  $W_R(M) \subseteq N_R(R)$
- b)  $J_R(R) \subseteq W_R(M)$  if  $M$  is a finitely generated  $R$ -module.

**Proof.** “(a)” Set  $x \in W_R(M)$ . Then  $xM \neq M$  and hence  $x$  is a non-unit element of  $R$ . Therefore  $x \in N_R(R)$ .

“(b)” Set  $x \in J_R(R)$ . Then  $Rx \subseteq J_R(R)$  and hence by the Nakayama lemma we have that  $xM = (Rx)M \neq M$ . □

**Theorem 3.3.** *Let  $M$  be an  $R$ -module. Then the following hold;*

- a)  $W_R(M) \subseteq N_R(M)$
- b) *We have equality in (a) if  $M$  is a finitely generated  $R$ -module.*

**Proof.** “(a)” If  $M = 0$  then there is nothing to prove. Let  $M \neq 0$  and let  $x \in W_R(M)$ . Then  $xM \neq M$  and hence  $M/xM \neq 0$ . Let  $\mathfrak{m} \in \text{MaxSupp}(M/xM)$ . Then  $(M/xM)_{\mathfrak{m}} \neq 0$  and hence  $M_{\mathfrak{m}}/(x/1)M_{\mathfrak{m}} \neq 0$ . Therefore by (2.2a) we have  $x/1 \in W_{R_{\mathfrak{m}}}(M_{\mathfrak{m}}) \subseteq N_{R_{\mathfrak{m}}}(R_{\mathfrak{m}}) = \mathfrak{m}R_{\mathfrak{m}}$  and hence  $x \in \mathfrak{m}$ . Now the assertion follows from the fact that  $\mathfrak{m} \in \text{MaxSupp}(M)$ .

“(b)” If  $M = 0$  then there is nothing to prove. Let  $M \neq 0$  and let  $x \in N_R(M)$ . Then there exists  $\mathfrak{m} \in \text{MaxSupp}(M)$  such that  $x \in \mathfrak{m}$ . Thus  $x/1 \in \mathfrak{m}R_{\mathfrak{m}} = J_{R_{\mathfrak{m}}}(R_{\mathfrak{m}})$ . Since  $M$  is a finitely generated  $R$ -module we have  $M_{\mathfrak{m}}$  is a non-zero finitely generated  $R_{\mathfrak{m}}$ -module. Therefore by (2.2b), we have that  $J_{R_{\mathfrak{m}}}(R_{\mathfrak{m}}) \subseteq W_{R_{\mathfrak{m}}}(M_{\mathfrak{m}})$  and hence  $x/1 \in W_{R_{\mathfrak{m}}}(M_{\mathfrak{m}})$ . Thus  $M_{\mathfrak{m}}/(x/1)M_{\mathfrak{m}} \neq 0$  and hence  $(M/xM)_{\mathfrak{m}} \neq 0$ . Therefore  $M/xM \neq 0$  and so  $x \in W_R(M)$ . □

**Theorem 3.4.** *Let  $M$  be an  $R$ -module. Then the following hold;*

a)  $Z_R(M) \subseteq N_R(M)$

b) *We have equality in (a) if  $M$  is a finitely cogenerated  $R$ -module.*

**Proof.** “(a)” Use [Y2; 1.1].

“(b)” Since  $M$  is finitely cogenerated we have  $\text{Supp}(M) \subseteq \text{MaxSpec}(R)$  and hence  $Z(M) = N(M)$  by [Y2; 1.1]. □

**Theorem 3.5.** *Let  $M$  be an injective  $R$ -module. Then the following hold;*

a)  $W_R(M) \subseteq Z_R(M)$

b) *We have equality in (a) if  $M$  is a finitely generated  $R$ -module.*

**Proof.** “(a)” Set  $x \in W_R(M)$ . If  $x \notin Z_R(M)$  we have the map  $\varphi : xM \rightarrow M$  with  $\varphi(xt) = t$  for any  $t \in M$ . Since  $M$  is an injective  $R$ -module, the map  $\varphi$  induces the map  $\psi : M \rightarrow M$  such that for all  $t \in M$  we have that  $t = \varphi(xt) = \psi(xt) = x\psi(t) \in xM$ . Thus  $xM = M$ , which is a contradiction.

“(b)” We have  $x \in Z_R(M) \subseteq N(M)$  by [Y2; 1.1]. Now the assertion follows from (3.3b). □

**Corollary 3.6.** If  $R$  is a self-injective ring then the set of zero divisors of  $R$  is equal to the set of non-units in  $R$ .

**Theorem 3.7.** *Let  $M$  be a flat  $R$ -module. Then the following hold;*

a)  $Z_R(M) \subseteq W_R(M)$

b) *We have equality in (a) if  $M$  is a finitely cogenerated  $R$ -module.*

**Proof.** “(a)” Set  $x \in Z_R(M)$ . Then there exists a non-zero element  $t \in M$  such that  $xt = 0$ . Thus we have the non-zero map  $\varphi : R/(x) \rightarrow M$  with  $\varphi(r + (x)) = rt$  for any  $r \in R$ . Therefore  $\text{Hom}(R/(x), M) \neq 0$  and hence there exists an injective module  $E$  such that  $\text{Hom}(\text{Hom}(R/(x), M), E) \neq 0$ . Since

$$\text{Hom}(\text{Hom}(R/(x), M), E) \cong R/(x) \otimes \text{Hom}(M, E),$$

we have that  $x \in W_R(\text{Hom}(M, E))$ . Since  $\text{Hom}(M, E)$  is an injective module we have  $x \in Z_R(\text{Hom}(M, E))$  by (3.4), and hence  $\text{Hom}(R/(x), \text{Hom}(M, E))$  is non-zero. Therefore  $\text{Hom}(R/(x) \otimes M, E) \neq 0$  and hence  $R/(x) \otimes M \neq 0$ . Thus  $x \in W_R(M)$

“(b)” By (3.3) and (3.4b) we have

$$W_R(M) \subseteq N_R(M) = Z_R(M).$$

Now the assertion follows from (a). □

## Acknowledgement

This paper was completed when the author visited (as an associate member) the International Centre for Theoretical Physics (ICTP), Trieste-Italy. The author wishes to thank Peyman Nasehpour and Keivan Mohajer, Shahid Beheshti University, for telling him about the (1.12) and (3.5). Also he wishes to thank Professor Chademan, University of Tehran, who suggested the problem about the set  $N'(R)$ . The author was supported in part by the University of Tehran and the Institute for studies in Theoretical Physics and Mathematics (IPM), Tehran-Iran.

## References

- [M] H. Matsumura, *Commutative ring theory*, Cambridge University Press, Cambridge 1986.
- [Y1] S. Yassemi, *Coassociated Primes*, Commun. in Algebra, **23** (1995), 1473–1498.
- [Y2] S. Yassemi, *Coassociated primes of modules over commutative rings*, Math. Scand., to appear.